

Module: General Preparedness

NNHS.Q173: Does your household have a 3-day supply of water for everyone who lives here? A 3-day supply of water is 1 gallon of water per person per day?

	Percent	Lower Estimate	Upper Estimate	N
Yes	77.3%	72.5%	81.4%	2,274
Male	76.9%	71.1%	81.7%	797
Female	76.2%	70.8%	80.9%	1,477

Does your household have a 3-day supply of water for everyone who lives here, by Age Group

	Point	Lower	Upper	N
<30	81.1%	73.3%	87.0%	317
30-39	61.6%	49.5%	72.4%	264
40-49	83.3%	75.3%	89.1%	326
50-59	78.1%	69.1%	85.1%	499
60-69	79.5%	69.3%	87.0%	447
70-79	84.5%	77.6%	89.5%	298
80+	66.6%	41.0%	85.1%	123

Does your household have a 3-day supply of water for everyone who lives here, by Agency

Agency	Point	Lower	Upper	N
Chinle	63.7%	53.7%	72.6%	698
Eastern	78.3%	72.5%	83.2%	467
Ft. Defiance	83.9%	79.3%	91.9%	339
Northern	65.7%	59.9%	71.2%	502
Western	81.5%	73.0%	87.8%	268

Data Source: Report of the Navajo Behavioral Risk Factor Surveillance Survey, 2013, 2015, 2016.

All Result Responses – General Preparedness.

Navajo Epidemiology Center/Navajo Department of Health

Page 1 of 6

Module: General Preparedness

NNHS.Q174: Does your household have a 3-day supply of nonperishable food for everyone who lives there? By nonperishable food we mean food that does not require refrigeration or cooking.

	Percent	Lower Estimate	Upper Estimate	N
Yes	82.8%	77.7%	87.0%	2,280
Male	82.8%	77.0%	87.3%	801
Female	82.8%	77.4%	87.1%	1,479

Does your household have a 3-day supply of nonperishable food for everyone who lives there, by Age Group

	Point	Lower	Upper	N
<30	79.9%	67.3%	88.4%	319
30-39	74.3%	62.1%	83.7%	263
40-49	87.9%	81.2%	92.5%	326
50-59	84.1%	74.9%	90.4%	498
60-69	87.9%	80.5%	92.8%	451
70-79	88.4%	81.9%	92.7%	300
80+	87.3%	78.3%	92.8%	123

Data Source: Report of the Navajo Behavioral Risk Factor Surveillance Survey, 2013, 2015, 2016.

All Result Responses – General Preparedness.

Navajo Epidemiology Center/Navajo Department of Health

Page 2 of 6

Module: General Preparedness

Does your household have a 3-day supply of nonperishable food for everyone who lives there, by Age Group

Agency	Point	Lower	Upper	N
Chinle	66.2%	55.2%	75.7%	698
Eastern	86.1%	80.5%	90.3%	467
Ft. Defiance	94.6%	90.0%	97.1%	337
Northern	75.0%	65.0%	82.8%	510
Western	84.8%	75.7%	90.8%	268

NNHS.Q175: Does your household have a 3-day supply of prescription medication for each person who takes prescribed medicine?

	Percent	Lower Estimate	Upper Estimate	N
Yes	83.3%	78.6%	87.2%	1,991
Male	83.1%	77.4%	87.6%	675
Female	83.1%	78.8%	86.7%	1,316

Module: General Preparedness

Does your household have a 3-day supply of prescription medication for each person who takes prescribed medicine, by Age Group

	Point	Lower	Upper	N
<30	77.4%	60.6%	88.4%	256
30-39	70.4%	58.9%	79.8%	215
40-49	89.1%	82.3%	93.4%	276
50-59	87.1%	77.8%	92.8%	436
60-69	92.9%	87.8%	95.9%	410
70-79	93.5%	89.6%	96.0%	281
80+	90.5%	73.2%	97.1%	117

Does your household have a 3-day supply of prescription medication for each person who takes prescribed medicine, by Agency

Agency	Point	Lower	Upper	N
Chinle	71.5%	67.0%	75.7%	612
Eastern	85.8%	78.3%	91.1%	390
Ft. Defiance	92.3%	87.5%	95.4%	319
Northern	88.1%	79.1%	93.5%	437
Western	75.2%	60.8%	85.6%	233

Data Source: Report of the Navajo Behavioral Risk Factor Surveillance Survey, 2013, 2015, 2016.

All Result Responses – General Preparedness.

Navajo Epidemiology Center/Navajo Department of Health

Page 4 of 6

Module: General Preparedness

NNHS.Q176: What would be your main method or way of getting information from authorities in a large-scale disaster or emergency? (N=2,279)

Method	Total	Male (N=797)	Female (1,482)
Television	23.2% (19.1%, 27.8%)	21.2% (16.4%, 27.1%)	24.4% (19.0%, 30.7%)
Radio	36.5% (32.0%, 41.1%)	41.3% (35.1%, 47.8%)	32.3% (27.7%, 37.2%)
Internet	8.6% (5.1%, 14.1%)	11.6% (6.2%, 20.7%)	6.1% (4.1%, 9.1%)
Print media	0.8% (0.3%, 2.5%)	0.9% (0.2%, 5.2%)	0.7% (0.3%, 1.6%)
Neighbors	10.6% (7.9%, 14.0%)	8.3% (5.4%, 12.6%)	12.6% (9.2%, 17.0%)
Other	14.4% (11.6%, 17.6%)	12.7% (8.6%, 18.3%)	15.8% (11.7%, 20.9%)
Don't know/Not sure	6.1% (2.9%, 12.2%)	3.9% (1.9%, 7.9%)	8.2% (3.3%, 19.2%)

Radio as the best method or way of getting information from authorities in a large-scale disaster or emergency, by Age Group

	Point	Lower	Upper	N
<30	33.6%	25.1%	43.2%	306
30-39	30.0%	19.8%	42.5%	253
40-49	34.8%	24.9%	46.2%	314
50-59	48.0%	39.6%	56.5%	483
60-69	40.1%	30.4%	50.7%	434
70-79	52.7%	42.0%	63.2%	293
80+	68.0%	51.5%	80.9%	117

Data Source: Report of the Navajo Behavioral Risk Factor Surveillance Survey, 2013, 2015, 2016.

All Result Responses – General Preparedness.

Navajo Epidemiology Center/Navajo Department of Health

Page 5 of 6

Module: General Preparedness

Radio as the best method or way of getting information from authorities in a large-scale disaster or emergency, by Agency

Agency	Point	Lower	Upper	N
Chinle	49.6%	45.2%	54.0%	685
Eastern	37.2%	25.7%	50.4%	456
Ft. Defiance	43.5%	36.6%	50.7%	321
Northern	28.3%	20.3%	37.9%	490
Western	35.5%	28.8%	42.8%	248

Note: 22.6% of those in the Northern agency said there was an “Other” method they would like to get disaster information, but no specific “Other” methods were identified.

